



## REFERENCES

1. Inouye SK, van Dyck CH, Alessi CA, et al. Clarifying confusion: the confusion assessment method. A new method for detection of delirium. *Ann Intern Med.* 1990;113:941-948.
2. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition.* 5th ed. Washington, DC: American Psychiatric Association; 2013.
3. Wong CL, Holroyd-Leduc J, Simel DL, Straus SE. Does this patient have delirium? *JAMA.* 2010;304:779-786.
4. Fick DM, Agostini JV, Inouye SK. Delirium superimposed on dementia: a systematic review. *J Am Geriatr Soc.* 2002;50:1723-1732.
5. Marcantonio ER, Goldman L, Mangione CM, et al. A clinical prediction rule for delirium after elective noncardiac surgery. *JAMA.* 1994;271:134-139.
6. Practice guideline for the treatment of patients with delirium. American Psychiatric Association. *Am J Psychiatry.* 1999;156(5 suppl):1-20.
7. Inouye SK. Delirium in older persons. *Engl J Med.* 2006;354:1157-1165.
8. Meagher DJ. Delirium: optimising management. *BMJ.* 2001;322:144-149.
9. Inouye SK, Charpentier PA. Precipitating factors for delirium in hospitalized elderly persons. Predictive model and interrelationship with baseline vulnerability. *JAMA.* 1996;275:852-857.
10. Elie M, Cole MG, Primeau FJ, Bellavance F. Delirium risk factors in elderly hospitalized patients. *J Gen Intern Med.* 1998;13:204-212.
11. Inouye SK, Bogardus ST, Jr, Charpentier PA, et al. A multicomponent intervention to prevent delirium in hospitalized older patients. *Engl J Med.* 1999;340:669-676.
12. Marcantonio ER, Flacker JM, Wright RJ, Resnick NM. Reducing delirium after hip fracture: a randomized trial. *J Am Geriatr Soc.* 2001;49:516-522.
13. Naughton BJ, Saltzman S, Ramadan F, et al. A multifactorial intervention to reduce prevalence of delirium and shorten hospital length of stay. *J Am Geriatr Soc.* 2005;53:18-23.
14. Michaud L, Bula C, Berney A, et al. Delirium: guidelines for general hospitals. *J Psychosom Res.* 2007;62:371-383.
15. Kalisvaart KJ, de Jonghe JF, Bogaards MJ, et al. Haloperidol prophylaxis for elderly hip surgery patients at risk for delirium: a randomized placebo-controlled study. *J Am Geriatr Soc.* 2005;53:1658-1666.
16. Roden DM. Drug-induced prolongation of the QT interval. *Engl J Med.* 2004;350:1013-1022.



17. Brigham and Women's Hospital Delirium Task Force. Acute Hospital Acquired Delirium Guidelines. Boston, MA; 2007.
18. Lacasse H, Perreault MM, Williamson DR. Systematic review of antipsychotics for the treatment of hospital-associated delirium in medically or surgically ill patients. *Ann Pharmacother*. 2006;40:1966-1973.
19. Ozbolt LB, Paniagua MA, Kaiser RM. Atypical antipsychotics for the treatment of delirious elders. *J Am Med Dir Assoc*. 2008;9:18-28.
20. Rea RS, Battistone S, Fong JJ, Devlin JW. Atypical antipsychotics versus haloperidol for treatment of delirium in acutely ill patients. *Pharmacotherapy*. 2007;27:588-594.
21. Lonergan E, Britton AM, Luxenberg J, Wyller T. Antipsychotics for delirium. *Cochrane Database Syst Rev*. 2007(2):CD005594.
22. Wang PS, Schneeweiss S, Avorn J, et al. Risk of death in elderly users of conventional vs. atypical antipsychotic medications. *N Engl J Med*. 2005;353:2335-2341.
23. Bradley KA, Rubinsky AD, Sun H, et al. Alcohol screening and risk of postoperative complications in male VA patients under-going major non-cardiac surgery. *J Gen Intern Med*. 2011;26:162-169.
24. Mayo-Smith MF, Beecher LH, Fischer TL, et al. Management of alcohol withdrawal delirium. An evidence-based practice guideline. *Arch Intern Med*. 2004;164:1405-1412.
25. Sullivan JT, Sykora K, Schneiderman J, et al. Assessment of alcohol withdrawal: the revised Clinical Institute Withdrawal Assessment for Alcohol scale (CIWA-Ar). *Br J Addict*. 1989;84:1353-1357.
26. Mayo-Smith MF. Pharmacological management of alcohol withdrawal. A meta-analysis and evidence-based practice guideline. American Society of Addiction Medicine Working Group on Pharmacological Management of Alcohol Withdrawal. *JAMA*. 1997;278:144-151.
27. Meagher DJ, et al. Concordance between DSM-IV and DSM V criteria for delirium diagnosis in a pooled database of 768 prospectively evaluated patients using delirium rating scale-revised-98. *BMC Medicine*. 2014;12:164
28. Fick DM, Inouye SK, Guess J, et al. Preliminary development of an ultrabrief two-item bedside test for delirium. *J Hosp Med*. 2015;10(00):000-000.
29. O'Regan NA, Ryan DJ, Boland E, et al. Attention! A good bedside test for delirium? *J Neurol Neurosurg Psychiatry*. 2014;85(10):1122-1131.
30. Inouye SK, Westendorp RG, Saczynski JS: Delirium in elderly people. *Lancet*. 2014, 383: 911-922
31. Watt D, Budding DE, Koziol LF. Delirium. Noggle CA, Dean RS (Eds.), *The neuropsychology of psychopathology*, Springer, NY (2013), 425-440.



32. Ryan DJ, O'Regan NA, Caoimh RÓ, Clare J, O'Connor M, Leonard M, McFarland J, Tighe S, O'Sullivan K, Trzepacz PT, Meagher D, Timmons S: Delirium in an adult acute hospital population: predictors, prevalence and detection. *BMJ Open*. 2013, 3: e001772-10.1136/bmjopen-2012-001772
33. Han JHm et al. Delirium in the emergency department: an independent predictor of death within 6 months. *Ann Emerg Med*. 2010;56:244-252
33. Buurman BM et al. Geriatric conditions in acutely hospitalized older patients: prevalence and one-year survival and functional decline. *PLoS One* 2011;6:e26951
34. Leslie DL, et al. Premature death associated with delirium at 1-year follow-up. *Arch Intern Med*. 2005;165:1657-62
35. Marcantonio ER, et al. Outcomes of older people admitted to postacute facilities with delirium. *J Am Geriatr Soc*. 2005;53:963-969.
36. Saczynski JS et al. Cognitive trajectories after postoperative delirium. *NEJM*. 2012;367:30-39.
37. Chedru F, Geschwind N. Disorders of higher cortical functions in acute confusional states. *Cortex* 1972;8:395-411
38. Jones RN, et al. Aging, brain disease, and reserve: implications for delirium. *Am J Geriatr Psychiatry*. 2010;18:117-127
39. Wilson RS, et al Cognitive decline after hospitalization in a community population of older persons. *Neurology*. 2012;78:950-956.
40. Saczynski JS, et al. Cognitive trajectories after postoperative delirium. 2012 *NEJM* 367:30-39
41. Cole MG, Primeau FJ. Prognosis of delirium in elderly hospital patients. *CMAJ*.1993;149(1):41
42. Gaudreau JD, Gagnon P. Psychotonic drugs and delirium pathogenesis: the central role of the thalamus. *Med Hypotheses* 2005;64:471-475
43. Leslie DL, et al. One-year health care costs associated with delirium in the elderly population. *Arch Intern Med* 2008;168:27-32
44. Wachter RM. *Understanding patient safety* (2nd edition), McGraw-Hill Medical, NY 2012.
45. Van Rompaey B, et al. The effect of earplugs during the night on the onset of delirium and sleep perception: a randomized controlled trial in intensive care patients. *Crit Care*. 2012 May;16(3):R73.



46. Leslie DL, et al. Consequences of preventing delirium in hospitalized older adults on nursing home costs. *J Am Geriatr Soc.* 2005;53:405-409
47. Campell N et al. Pharmacological management of delirium in hospitalized adults – a systematic evidence review. *J Gen Intern Med.* 2009;24(7):848.
48. Atalan N, et al. Morphine is a reasonable alternative to haloperidol in the treatment of postoperative hyperactive-type delirium after cardiac surgery. *J Cardiothorac Vasc Anesth.* 2013 Oct;27(5):933-38.
49. Moryl N, Comfort C, Obbens E. Methadone in the treatment of pain and terminal delirium in advanced cancer patients. *Palliat Support Care.* 2005;3(4):311.
50. Hudetz JA, et al. Ketamine attenuates delirium after cardiac surgery with cardiopulmonary bypass. *J Cardiothorac Vasc Anesth.* 2009 Oct;23(5):651-657.)
51. Reade MC et al. Effect of dexmedetomidine added to standard care on ventilator-free time in patients with agitated delirium: A randomized clinical trial. *JAMA* 2016 Apr 12; 315:1460.
52. Awissi DK, et al. Alcohol withdrawal and delirium tremens in the critically ill: a systematic review and commentary. *Intensive Care Med.* 2013 Jan;39(1):16-30.
53. Saitz R, Mayo-Smith MF, Roberts MS, Redmond HA, Bernard DR, Calkins DR. Individualized treatment for alcohol withdrawal. a randomized double-blind controlled trial. *JAMA.* 1994;272:519–23.
54. Daeppen JB, Gache P, Landry U, Sekera E, Schweizer V, Gloor S, et al. Symptom-triggered vs fixed-schedule doses of benzodiazepine for alcohol withdrawal: a randomized treatment trial. *Arch Intern Med.* 2002;162:1117–21.
55. Cassidy EM, et al. Symptom-triggered benzodiazepine therapy for alcohol withdrawal syndrome in the emergency department: a comparison with the standard fixed dose benzodiazepine regimen. *Emerg Med J.* 2012 Oct;29(10):802-4.
56. Sullivan JT, Swift RM, Lewis DC. Benzodiazepine requirements during alcohol withdrawal syndrome: clinical implications of using a standardized withdrawal scale. *J Clin Psychopharmacol.* 1991;11:291–95.
57. Reoux JP, Miller K. Routine hospital alcohol detoxification practice compared with symptom triggered management with an Objective Withdrawal Scale (CIWA-Ar) *Am J Addict.* 2000;9:135–44.
58. Jaeger T, Lohr R, Pankratz VS. Symptom-triggered therapy for alcohol withdrawal syndrome in medical inpatients. *Mayo Clinic Procedure.* 2001;77:695–701.
59. Ferguson JA, et al. Risk factors for delirium tremens development. *J Gen Intern Med* 1996;11(7):410.



60. Monte R , et al. Risk factors for delirium tremens in patients with alcohol withdrawal syndrome in a hospital setting. *Eur J Intern Med.* 2009 Nov;20(7):690-4
61. Kim DW, et al. Clinical predictors for delirium tremens in patients with alcohol withdrawal seizures. *Am J Emerg Med.* 2015 May;33(5):701-4
62. Won A, Smithburger PL, Kane-Gill SL. Review of adjunctive dexmedetomidine in the management of severe acute alcohol withdrawal syndrome. *Am J Drug Alcohol Abus.* 2015;41(5):382-91.
63. Malcolm R, Ballenger JC, Sturgis ET, Anton R. Double-blind controlled trial comparing carbamazepine to oxazepam treatment of alcohol withdrawal. *Am J Psychiatry.* 1989;146:617–21.
64. Chourishi A, Raichandani OP, Chandraker S, Chourishi S. A comparative study of efficacy and tolerability of lorazepam and gabapentin in the treatment of alcohol withdrawal syndrome. *IJPSR.* 2010;3(2):80–84.
65. Stuppaeck CH, Deisenhammer EA, Kurz M, Whitworth AB, Hinterhuber H. The irreversible gamma-aminobutyrate transaminase inhibitor vigabatrin in the treatment of the alcohol withdrawal syndrome. *Alcohol Alcohol.* 1996;31:109–11.
66. Mirijello A et al. Identification and management of alcohol withdrawal syndrome. *Drugs.* 2015 Mar;75(4):353-65.
67. Wei LA, et al. The confusion assessment method: a systematic review of current usage. *J Am Geriatr Soc.* 2008;56:823-830