

Management of Patients with Hip Fractures

(2019-2022)

References

1. Samelson EJ, Zhang Y, Kiel DP, et al. Effect of birth cohort on risk of hip fracture: age-specific incidence rates in the Framingham Study. *Am J Public Health*. 2002;92:858-862.
2. Centers for Disease Control. Home and Recreational Safety Important Facts about Falls. 2017 Feb 10. <https://www.cdc.gov/homeandrecreationalafety/falls/adultfalls.html>.
3. Bhandari M, Swiontkowski M. Management of Acute Hip Fracture. *N Engl J Med*. 2017;377:2053-2062.
4. Cauley JA. Osteoporosis: fracture epidemiology update 2016. *Curr Opin Rheumatol*. 2017;29:150-156.
5. DeFrances CJ, Cullen KA, Kozak LJ. National Discharge Survey: 2005 annual summary with detailed diagnosis and procedure data. National Center for Health Statistics. *Vital Health Stat 13*. 2007;(165):1-209.
6. Zuckerman JD. Hip fracture. *N Engl J Med*. 1996;334:1519-1525.
7. Leibson CL, Tosteson ANA, Gabriel SE, et al. Mortality, disability, and nursing home use for persons with and without hip fracture: a population-based study. *J Am Geriatr Soc*. 2002;50:1644-1650.
8. Magaziner J, Hawkes W, Hebel JR, et al. Recovery from hip fracture in eight areas of function. *J Gerontol A Biol Sci Med Sci*. 2000;55:M498-M507.
9. Veronese N, Maggi S. Epidemiology and social costs of hip fracture. *Injury*. 2018 Apr. [Epub ahead of print].
10. Johnston YA, Bergen G, Bauer M, et al. Implementation of the Stopping Elderly Accidents, Deaths, and Injuries Initiative in Primary Care: An Outcome Evaluation. *Gerontologist*. 2018 Sep. gny101.
11. Gillespie WJ, Gillespie LD, Parker MJ. Hip protectors for preventing hip fractures in older people. *Cochrane Database Syst Rev*. 2010;(10):CD001255.
12. Sharma G, Kuo Y-F, Freeman J, et al. Comanagement of hospitalized surgical patients by medicine physicians in the United States. *Arch Intern Med*. 2010 Feb 22;170(4):363-368.
13. Roy A, Heckman MG, Roy V. Associations between the hospitalist model of care and quality-of-care-related outcomes in patients undergoing hip fracture surgery. *Mayo Clin Proc*. 2006;81:28-31.
14. Swart E, Kates S, McGee S, et al. The Case for Comanagement and Care Pathways for Osteoporotic Patients with a Hip Fracture. *J Bone Joint Surg Am*. 2018;100:1343-50.
15. Swart E, Vasudeva BS, Makhni EC, et al. Dedicated perioperative hip fracture comanagement programs are cost-effective in high-volume centers: An economic analysis. *Clin Orthop Relat Res*. 2016 Jan;474(1):222-223.
16. American Academy of Orthopaedic Surgeons. Management of Hip Fractures in the Elderly. Evidence-Based Clinical Practice Guidelines. September 5, 2014. <https://www.aaos.org/CustomTemplates/Content.aspx?id=6395&ssopc=1>
17. Hoffman R, Haas NP. Trochanteric Fractures (31-A) AO Principles of Fracture Management. AO Foundation Publishing. https://www2.aofoundation.org/wps/portal/!ut/p/a0/04_Sj9CPykssy0xPLMnMz0vMAfGjzOKN_A0M3D2_DDdbz9_UMMDRyDXQ3dw9wMDAx8jfULsh0VAdAsNSU!/?bone=Femur&segment=Proximal&soloState=lyteframe&contentUrl=srg/popup/further_reading/PFxM2/31/661_22-troc_fxs_surg_treatm.jsp. Accessed Feb 7, 2019.
18. Pincus D, Ravi B, Wasserstein D, et al. Association between wait time and 30-day mortality in adults undergoing hip fracture surgery. *JAMA*. 2017 Nov 28;318(20):1994-2003.
19. Fu MC, Boddapati V, Gausden EB, et al. Surgery for a fracture of the hip within 24 hours of admission is independently associated with reduced short-term post-operative complications. *Bone Joint J*. 2017 Sep;99-B(9):1216-1222.
20. Fleisher LA, Fleischmann KE, Auerbach AD, et al. 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2014;64:e77-137.
21. Grant PJ, Wesorick DH. Perioperative medicine for the hospitalized patient. *Med Clin N Am*. 2008;92:325-348.

22. Ellenbogen MI, Brotman DJ, Prichett L, et al. Contemporary rates of preoperative cardiac testing prior to inpatient hip fracture surgery. *J Hosp Med (Online)*. 2019; **14**(4): 224-8.
23. Lee TH, Marcantonio ER, Mangione CM, et al. Derivation and prospective validation of a simple index for prediction of cardiac risk of major noncardiac surgery. *Circulation*. 1999;100:1043-1049.
24. Whinney CM. Do hip fractures need to be repaired within 24 hours of injury? *Cleve Clin J Med*. 2005;72:250-252.
25. McFalls EO, Ward HB, Moritz TE, et al. Coronary artery revascularization before elective major vascular surgery. *N Engl J Med*. 2004;351:2795-804.
26. Stitgen A, Poludnianyk K, Dulaney-Crip E, et al. Adherence to preoperative cardiac clearance guidelines in hip fracture patients. *J Orthop Trauma*. 2015 Nov;29(11):500-3.
27. Lin H-S, Watts JN, Peel NM, et al. Frailty and post-operative outcomes in older surgical patients: a systematic review. *BMC Geriatr*. 2016 Aug 31;16(1):157.
28. Rockwood K, Song X, MacKnight C, et al. A global clinical measure of fitness and frailty in elderly people. *CMAJ*. 2005;173(5):489-95.
29. McIsaac DI, Taljaard M, Bryson GL, et al. Frailty as a predictor of death or new disability after surgery: A prospective cohort study. *Ann Surg*. 2018; Jul 24. doi: 10.1097/SLA.
30. Belga S, Majumdar SR, Kahlon S, et al. Comparing three different measures of frailty in medical inpatients: Multicenter prospective cohort study examining 30-day risk of readmission or death. *J Hosp Med*. 2016. August;11(8):556-562.
31. Berry SD, Rothbaum, RR, Kiel DP, et al. Association and clinical outcomes with surgical repair of hip fracture vs nonsurgical management in nursing home residents with advanced dementia. *JAMA Intern Med*. 2018;178(6):774-780.
32. Bohl DD, Shen MR, Hannon CP, et al. Serum albumin predicts survival and postoperative course following surgery for geriatric hip fracture. *J Bone Joint Surg Am*. 2017 Dec 20;99(24):2110-2118.
33. Taylor BE, McClave SA, Martindale RG, et al. Guidelines for the provision and assessment of nutrition support therapy in the adult critically ill patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.). *Crit Care Med*. 2016 Feb;44(2):390-438.
34. Geerts WH, Bergqvist D, Pineo GF, et al. Prevention of venous thromboembolism. American College of Chest Physicians Evidence-Based Clinical Practice Guidelines, 8th ed. *Chest*. 2008;133:381S-453S.
35. Falck-Ytter Y, Francis CW, Johanson NA, et al. Prevention of VTE in Orthopedic Surgery Patients. Antithrombotic Therapy and Prevention of Thrombosis, 9th ed. American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest*. 2012;141(2)(suppl):e278S-e325S.
36. Balk EM, Ellis AG, Di M, Adam GP, Trikalinos TA. Venous Thromboembolism Prophylaxis in Major Orthopedic Surgery: Systematic Review Update. Comparative Effectiveness Review No. 191. (Prepared by the Brown Evidence-based Practice Center under Contract No. 290-2015-00002-I.) AHRQ Publication No. 17-EHC021-EF. Rockville, MD: Agency for Healthcare Research and Quality; June 2017. www.effectivehealthcare.ahrq.gov/reports/final.cfm.
37. Eriksson BI, Bauer KA, Lassen MR, et al. Fondaparinux compared with enoxaparin for the prevention of venous thromboembolism after hip-fracture surgery. *N Engl J Med*. 2001;345:1298-304.
38. Fuji T, Fujita S, Kawai Y, et al. Safety and efficacy of edoxaban in patients undergoing hip fracture surgery. *Thrombosis Research*. 2014 June;133(6):1016-1022.
39. Sasaki S, Miyakoshi N, Matsuura H, et al. Prospective study on the efficacies of fondaparinux and enoxaparin in preventing venous thromboembolism after hip fracture surgery. *J Orthop Sci*. 2011;16: 64-70.
40. Anderson DR, Morgano GP, Bennett C, et al. American Society of Hematology 2019 guidelines for management of venous thromboembolism: prevention of venous thromboembolism in surgical hospitalized patients. *Blood Adv*. 2019;10:3(23):3898-3944
41. Wong A, Kraus PS, Lau BD, et al. Patient preferences regarding pharmacologic venous thromboembolism prophylaxis. *J Hosp Med*. 2015 Feb;10(2):108-11.
42. Drescher FS, Sirovich BE, Lee A, et al. Aspirin versus Anticoagulation for Prevention of Venous Thromboembolism Major Lower Extremity Orthopedic Surgery: A Systematic Review and Meta-Analysis. *J Hosp Med*. 2014;9:579-585.
43. Guyatt GH, Cook DJ, Jaeschke R, et al. Grades of recommendation for antithrombotic agents: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines, 8th ed. *Chest*. 2008;133:123S-131S.

44. Hung WW, Egol KA, Zuckerman JD, Siu AL. Hip fracture management tailoring Care for the older patient. *JAMA*. 2012;307(20):2185-2194.
45. Yang Y, Zhao X, Dong T, et al. Risk factors for postoperative delirium following hip fracture repair in elderly patients: a systematic review and meta-analysis. *Aging Clin Exp Res*. 2017 Apr;29(2):115-126.
46. Eamer G, Taheri A, Chen SS, et al. Comprehensive geriatric assessment for older people admitted to a surgical service. *Cochrane Database Syst Rev*. 2018 Jan 31;1:CD012485.
47. Shields L, Henderson V, Caslake R. Comprehensive geriatric assessment for prevention of delirium after hip fracture: a systematic review of randomized controlled trials. *J Am Geriatr Soc*. 2017 Jul;65(7):1559-1565.
48. The American Geriatrics Society Expert Panel on Postoperative Delirium in Older Adults. American Geriatrics Society Abstracted Clinical Practice Guideline for Postoperative Delirium in Older Adults. *J Am Geriatr Soc*. 2015 January;63(1):142-150.
49. Guay J, Parker MJ, Griffiths R, Kopp S. Peripheral nerve blocks for hip fractures. *Cochrane Database Syst Rev*. 2017 May 11;5:CD001159.
50. Beaupre LA, Jones, CA, Saunders LD, et al. Best practices for elderly hip fracture patients. A systematic overview of the evidence. *J Gen Intern Med*. 2005;20:09-1025.
51. Baumgarten M, Margolis D, Berlin JA, et al. Risk factors for pressure ulcers among elderly hip fracture patients. *Wound Rep Reg*. 2003;11:96-103.
52. Hofman A, Geelkerken RH, Wille J, et al. Pressure sores and pressure-decreasing mattresses: controlled clinical trial. *Lancet*. 1994;343:568-571.
53. Siris ES, Adler R, Bilezikian J, et al. The clinical diagnosis of osteoporosis: a position statement from the National Bone Health Alliance Working Group. *Osteoporos Int*. 2014: 1439-1443.
54. Colon-Emeric C, Nordsletten L, Olson S, et al. Association between timing of zoledronic acid infusion and hip fracture healing. *Osteoporosis Int*. 2011;22:2329-2336.
55. Cosman F, de Beur SJ, LeBoff MS, et al. Clinician's Guide to Prevention and Treatment of Osteoporosis. *Osteoporosis Int*. 2014 Aug.
56. US Preventive Services Task Force. Vitamin D, calcium, or combined supplementation for the primary prevention of fractures in community-dwelling adults. US Preventive Services Task Force Recommendation Statement. *JAMA*. 2018;319(15):1592-1599.
57. Bolland MJ, Grey A, Avenell A. Effects of vitamin D supplementation on musculoskeletal health: a systematic review, meta-analysis, and trial sequential analysis. *Lancet Diabetes Endocrinol*. 2018 November;6(11):847-858.
58. Lyles KW, Colon-Emeric CS, Magaziner JS, et al. Zoledronic Acid in Reducing Clinical Fracture and Mortality after Hip Fracture. *N Engl J Med*. 2007;357:nihpa40967.
59. Li YT, Cai HF, Zhang ZL. Timing of the initiation of bisphosphonates after surgery for fracture healing: a systematic review and meta-analysis of randomized controlled trials. *Osteoporosis Int*. 2015;26:431-441.
60. Unnanuntana A, Laohaprasitiporn P, Jarusriwanna A. Effect of bisphosphonate initiation at week 2 versus week 12 on short-term functional recovery after femoral neck fracture: a randomized controlled trial. *Arch Osteoporos*. 2017 Dec;12(1):27.
61. Desai RJ, Mahesri M, Abdia Y, et al. Association of Osteoporosis Medication Use after Hip Fracture with Prevention of Subsequent Nonvertebral Fractures An Instrumental Variable Analysis. *JAMA Network Open*. 2018;1(3):e180826.
62. Tan EK, Loh KP, Goff SL. Internal Medicine Hospitalists' Perceived Barriers and Recommendations for Optimizing Secondary Prevention of Osteoporotic Hip Fractures. *South Med J*. 2017 Dec;110(12):749-753.
63. Lv H, Yin P, Long A, et al. Clinical characteristics and risk factors of postoperative pneumonia after hip fracture surgery: a prospective cohort study. *Osteoporos Int*. 2016 Oct;27(10): 3001-9.