

References

1. Esteban NV, Loughlin T, Yergey AL, et al. Daily cortisol production rate in men determined by stable isotope dilution/mass spectrometry. *J Clin Endocr Metab.* 1991;72:39-44.
2. Loriaux DL. Adrenal. In: Nabel EG, ed. *ACP Medicine.* Hamilton, Ontario, Canada: BC Decker; 2009:3 Endo IV, 1-13.
3. Eisenbarth GS, Gottleib PA. Autoimmune polyendocrine syndromes. *N Engl J Med.* 2004;350:2068-2079.
4. Brunton LL, ed. *Goodman & Gilman's The Pharmacologic Basis of Therapeutics.* 12th ed. New York, NY: McGraw-Hill; 2011:1216.
5. Schiff RL, Welsh GA. Perioperative evaluation and management of the patient with endocrine dysfunction. *Med Clin N Am.* 2003;87:175-192.
6. Krasner AS. Glucocorticoid-induced adrenal insufficiency. *JAMA.* 1999;282:671-676.7.
Dangodara AA, Muluk VS. Adrenal insufficiency. In Cohn SL (editor). *Perioperative Medicine.* New York, NY; Springer-Verlag; 2011;235-242.
8. Marik PE, Varon J. Requirement of perioperative stress doses of corticosteroids; a systematic review of the literature. *Arch Surg.* 2008;143:1222-1226
9. Gilbert K, Lewis JB Jr, Smetana GW. Assessing perioperative risk. In: Cohn SL, Smetana GW, Weed HG, eds. *Perioperative Medicine: Just the Facts.* New York, NY: McGraw-Hill; 2006:9-17.
10. MacKenzie CR, Goodman SM. Stress dose steroids: myths and perioperative medicine. *Curr Rheumatol Rep;* 2016;18:47.