

Managing Diabetes and Hyperglycemia in the Hospital:

Critically Ill and Surgical Patients

(2023-2026)

REFERENCES

1. Clement S, Baithwaite SS, Magee MF, et al. Management of diabetes and hyperglycemia in hospitals. *Diabetes Care*. 2004;27:553-591.
2. Van den Berghe G, Wouters P, Weekers F, et al. Intensive insulin therapy in critically ill patients. *N Engl J Med*. 2001;345:1359-1367.
3. Van den Berghe G, Wilmer A, Hermans G, et al. Intensive insulin therapy in the medical ICU. *N Engl J Med*. 2006;354:449-461.
4. Brunkhorst FM, Engel C, Bloos F, et al. Intensive insulin therapy and pentastarch resuscitation in severe sepsis. *N Engl J Med*. 2008;358:125-139.
5. Wiener RS, Weiner DC, Larson RJ. Benefits and risks of tight glucose control in critically ill adults: a meta-analysis. *JAMA*. 2008;300:933-944.
6. The NICE-SUGAR Study Investigators, Finfer S, Chittock DR, et al. Intensive versus conventional glucose control in critically ill patients. *N Engl J Med*. 2009;360:1283-1297.
7. Inzucchi SE, Siegel MD. Glucose control in the ICU-how tight is too tight? *N Engl J Med*. 2009;360:1346-1349.
8. Griesdale DEG, de Souza RJ, van Dam RM, et al. Intensive insulin therapy and mortality among critically ill patients: a meta-analysis including NICE-SUGAR study data. *CMAJ*. 2009;180:821-827.
9. Qaseem A, Humphrey LL, Chou R, et al. Use of intensive insulin therapy for the management of glycemic control in hospitalized patients: a clinical practice guideline from the American College of Physicians. *Ann Intern Med*. 2011;154:260-267.
10. Lipshutz AK, Gropper MA. Perioperative glycemic control: an evidence-based review. *Anesthesiology*. 2009;110:408-421.
11. Dronge AS, Perkal MF, Kancir S, et al. Long-term glycemic control and postoperative infectious complications. *Arch Surg*. 2006;141:375-380; discussion 380.
12. Vann MA. Perioperative management of ambulatory surgical patients with diabetes mellitus. *Curr Opin Anaesthesiol*. 2009;22:718-724.

13. Lazar HL, McDonnell M, Chipkin SR, et al. The Society of Thoracic Surgeons practice guideline series: blood glucose management during adult cardiac surgery. *Ann Thorac Surg.* 2009;87:663-669.
14. Furnary A, Gao G, Grunkemeier G, et al. Continuous insulin infusion reduces mortality in patients with diabetes undergoing coronary artery bypass grafting. *J Thorac Cardiovasc Surg.* 2003;125:1007-1021.
15. Furnary A, Wu Y, Bookin S. Effect of hyperglycemia and continuous intravenous insulin infusions on outcomes of cardiac surgical procedures: the Portland Diabetic Project. *Endocr Pract.* 2004;10 (suppl 2):21-33.
16. Lazar HL, Chipkin SR, Fitzgerald CA, et al. Tight glycaemic control in diabetic coronary artery bypass graft patients improves perioperative outcomes and decreases recurrent ischemic events. *Circulation.* 2004;109:1497-1502.
17. Doenst T, Wijeyesundera D, Karkouti K, et al. Hyperglycemia during cardiopulmonary bypass is an independent risk factor for mortality in patients undergoing cardiac surgery. *J Thorac Cardiovasc Surg.* 2005;130:1144.
18. Ouattara A, Lecomte P, Le Manach Y, et al. Poor intraoperative blood glucose control is associated with a worsened hospital outcome after cardiac surgery in diabetic patients. *Anesthesiology.* 2005;103:687-694.
19. Gandhi GY, Nuttall GA, Abel MD, et al. Intraoperative hyperglycemia and perioperative outcomes in cardiac surgery patients. *Mayo Clin Proc.* 2005;80:862-866.
20. Subramaniam B, Panzica PJ, Novack V, et al. Continuous perioperative insulin infusion decreases major cardiovascular events in patients undergoing vascular surgery: a prospective, randomized trial. *Anesthesiology.* 2009;110:970-977.
21. Gandhi GY, Nuttall GA, Abel MD, et al. Intensive intraoperative insulin therapy versus conventional glucose management during cardiac surgery: a randomized trial. *Ann Intern Med.* 2007;146:233-243.
22. Moghissi E, Korytkowski M, DiNardo M, et al. American Association of Clinical Endocrinologists and American Diabetes Association consensus statement on inpatient glycaemic control. *Endocr Pract.* 2009;15:353-369.
23. Sourij H, Schmölder I, Kettler-Schmut E, et al. Efficacy of a continuous GLP-1 infusion compared with a structured insulin infusion protocol to reach normoglycemia in nonfasted type 2 diabetic patients: a clinical pilot trial. *Diabetes Care.* 2009;32:1669-1671.
24. Deane AM, Chapman MJ, Fraser RJ, et al. The effect of exogenous glucagon-like peptide-1 on the glycaemic response to small intestinal nutrient in the critically ill: a randomised double-blind placebo-controlled cross over study. *Crit Care.* 2009;13:R67.

25. Meier JJ, Weyhe D, Michaely M, et al. Intravenous glucagon-like peptide 1 normalizes blood glucose after major surgery in patients with type 2 diabetes. *Crit Care Med*. 2004;32:848-851.
26. Sokos GG, Bolukoglu H, German J, et al. Effect of glucagon-like peptide-1 (GLP-1) on glycemic control and left ventricular function in patients undergoing coronary artery bypass grafting. *Am J Cardiol*. 2007;100:824-829.
27. Schwartz S, DeFronzo RA. Is incretin-based therapy ready for the care of hospitalized patients with type 2 diabetes?: The time has come for GLP-1 receptor agonists! *Diabetes Care*. 2013;36:2107-2111.
28. Czosnowski QA, Swanson JM, Lobo BL, et al. Evaluation of glycemic control following discontinuation of an intensive insulin protocol. *J Hosp Med*. 2009;4:28-34.
29. Ramos P, Childers D, Maynard G, et al. Maintaining glycemic control when transitioning from infusion insulin: a protocol-driven, multidisciplinary approach. *J Hosp Med*. 2010;5:446-451.
30. Lepore M, Pampanelli S, Fanelli C, et al. Pharmacokinetics and pharmacodynamics of subcutaneous injection of long-acting human insulin analog glargine, NPH insulin, and ultralente human insulin and continuous subcutaneous infusion of insulin lispro. *Diabetes*. 2000;49:2142-2148.
31. Schmeltz LR, DeSantis AJ, Schmidt K, et al. Conversion of intravenous insulin infusions to subcutaneously administered insulin glargine in patients with hyperglycemia. *Endocr Pract*. 2006;12:641-650.
32. Bode BW, Braithwaite SS, Steed RD, Davidson PC. Intravenous insulin infusion therapy: indications, methods, and transition to subcutaneous insulin therapy. *Endocr Pract*. 2004;10(suppl 2):71-80.
33. DeSantis AJ, Schmeltz LR, Schmidt K, et al. Inpatient management of hyperglycemia: the Northwestern experience. *Endocr Pract*. 2006;12:491-505.
34. Furnary AP, Braithwaite SS. Effects of outcome on in-hospital transition from intravenous insulin infusion to subcutaneous therapy. *Am J Cardiol*. 2006;98:557-564.
35. LaPar et al. *J Thorac Cardiovasc Surg* 2014 Mar;147(3):1041-8. Joint Commission Online http://www.jointcommission.org/assets/1/23/jconline_January_28_15.pdf
36. American Diabetes Association. 15. Diabetes Care in the Hospital: *Standards of Medical Care in Diabetes-2021*. *Diabetes Care*. 2021 Jan;44(Suppl 1):S211-S220. doi: 10.2337/dc21-S015.
37. Blonde L, Umpierrez GE, Reddy SS, McGill JB, Berga SL, Bush M, Chandrasekaran S, DeFronzo RA, Einhorn D, Galindo RJ, Gardner TW, Garg R, Garvey WT, Hirsch IB, Hurley DL, Izuora K, Kosiborod M, Olson D, Patel SB, Pop-Busui R, Sadhu AR, Samson SL, Stec C, Tamborlane WV Jr, Tuttle KR, Twining C, Vella A, Vellanki P, Weber SL. American Association of Clinical Endocrinology Clinical Practice Guideline: Developing a Diabetes Mellitus Comprehensive Care Plan-2022 Update. *Endocr Pract*. 2022 Oct;28(10):923-1049. doi: 10.1016/j.eprac.2022.08.002.

38. Qaseem A, Humphrey LL, Chou R, Snow V, Shekelle P; Clinical Guidelines Committee of the American College of Physicians. Use of intensive insulin therapy for the management of glycemic control in hospitalized patients: a clinical practice guideline from the American College of Physicians. *Ann Intern Med.* 2011 Feb 15;154(4):260-7. doi: 10.7326/0003-4819-154-4-201102150-00007.

39. Lazar HL, McDonnell M, Chipkin SR, Furnary AP, Engelman RM, Sadhu AR, Bridges CR, Haan CK, Svedjeholm R, Taegtmeier H, Shemin RJ; Society of Thoracic Surgeons Blood Glucose Guideline Task Force. The Society of Thoracic Surgeons practice guideline series: Blood glucose management during adult cardiac surgery. *Ann Thorac Surg.* 2009 Feb;87(2):663-9. doi: 10.1016/j.athoracsur.2008.11.011.

40. Giori, NJ, Ellerbe LS, Bowe T, et al. Many Diabetic Total Joint Arthroplasty Candidates Are Unable to Achieve a Preoperative Hemoglobin A1c Goal of 7% or Less. *J Bone Jt Surg* 2014;96(6):500-504.