

Management of Patients with Hip Fracture

(2023-2026)

REFERENCES

1. Bhandari M, Swiontkowski M. Management of Acute Hip Fracture. *N Engl J Med*. 2017;377:2053-2062.
2. Samelson EJ, Zhang Y, Kiel DP, et al. Effect of birth cohort on risk of hip fracture: age-specific incidence rates in the Framingham Study. *Am J Public Health*. 2002;92:858-862.
3. HCUPnet. Healthcare Cost and Utilization Project (HCUP). 2012. Agency for Healthcare Research and Quality, Rockville, MD. <http://hcupnet.ahrq.gov>
4. Centers for Disease Control. Home and Recreational Safety Important Facts about Falls. 2017 Feb 10. <https://www.cdc.gov/homeandrecreationalafety/falls/adultfalls.html>.
5. GBD 2019 Fracture Collaborators. Global, regional, and national burden of bone fractures in 204 countries and territories, 1990–2019: a systematic analysis from the Global Burden of Disease Study 2019. *Lancet Health Long*. 2021;2(9):e580-e592.
6. Williams SA, Chastek B, Sundquist K, et al. Economic burden of osteoporotic fractures in US managed care enrollees. *Am J Manag Care*. 2020;26(5):e142-e149.
7. Remily EA, Mohamed NS, Wilkie WA, et al. Hip Fracture Trends in America Between 2009 and 2016. *Geriatr Ortho Surg Rehab*. 2020; 11. doi:10.1177/2151459320929581.
8. Harvey N, Dennison E, Cooper C. Osteoporosis: impact on health and outcomes. *Nat Rev Rheumatol*. 2010; 6:99–105.
9. Cauley JA. Osteoporosis: fracture epidemiology update 2016. *Curr Opin Rheumatol*. 2017;29:150-156.
10. Tsang C, Bolton C, Burgon V, et al. Predicting 30 day mortality after hip fracture: evaluation of the national hip fracture database case-mix adjustment model. *Bone Joint Res*. 2017;5(9):550-556.
11. Downey C, Kelly M, Quinlan JF. Changing trends in the mortality rate at 1-year post hip fracture - a systematic review. *World J Orthop*. 2019;10(3):166-175.
12. Zuckerman JD. Hip fracture. *N Engl J Med*. 1996;334:1519-1525.

13. Leibson CL, Tosteson ANA, Gabriel SE, et al. Mortality, disability, and nursing home use for persons with and without hip fracture: a population-based study. *J Am Geriatr Soc*. 2002;50:1644-1650.
14. Dyer SM, Crotty M, Fairhall N, et al. A critical review of the long-term disability outcomes following hip fracture. *BMC Geriatr*. 2016;16(1):158.
15. Wright NC, Saag KG, Curtis JR, et al. Recent trends in hip fracture rates by race/ethnicity among older US adults. *J Bone Miner Res*. 2012;27:2325-2332.
16. Thorell K, Ranstad K, Midlöv P et al. Is use of fall risk-increasing drugs in an elderly population associated with an increased risk of hip fracture, after adjustment for multimorbidity level: a cohort study. *BMC Geriatr*. 2014;4(14):131.
17. Johnston YA, Bergen G, Bauer M, et al. Implementation of the Stopping Elderly Accidents, Deaths, and Injuries Initiative in Primary Care: An Outcome Evaluation. *Gerontologist*. 2019;59(6):1182-1191.
18. Veronese N, Maggi S. Epidemiology and social costs of hip fracture. *Injury*. 2018;49(8):1458-1460.
19. Gillespie WJ, Gillespie LD, Parker MJ. Hip protectors for preventing hip fractures in older people. *Cochrane Database Syst Rev*. 2010;(10):CD001255.
20. Sharma G, Kuo Y-F, Freeman J, et al. Comanagement of hospitalized surgical patients by medicine physicians in the United States. *Arch Intern Med*. 2010;170(4):363-368.
21. Rohatgi N, Weng Y, Kittle J, Ahuja N. Merits of Surgical Comanagement of Patients With Hip Fracture by Dedicated Orthopaedic Hospitalists. *J Am Acad Orthop Surg Glob Res Rev*. 2021;5(3):e20.00231.
22. Roy A, Heckman MG, Roy V. Associations between the hospitalist model of care and quality-of-care-related outcomes in patients undergoing hip fracture surgery. *Mayo Clin Proc*. 2006;81:28-31.
23. Swart E, Kates S, McGee S, et al. The Case for Comanagement and Care Pathways for Osteoporotic Patients with a Hip Fracture. *J Bone Joint Surg Am*. 2018;100:1343-50.
24. Maxwell BG, Mirza A. Medical Comanagement of Hip Fracture Patients Is Not Associated with Superior Perioperative Outcomes: A Propensity Score-Matched Retrospective Cohort Analysis of the National Surgical Quality Improvement Project. *J Hosp Med*. 2020;15(8):468-474.
25. Batsis JA, Phy MP, Melton LJ, et al. Effects of a hospitalist care model on mortality of elderly patients with hip fractures. *J Hosp Med*. 2007;2(4):219-225.

26. Mazzarello S, McIsaac DI, Montroy J, et al. Postoperative shared-care for patients undergoing non-cardiac surgery: a systematic review and meta-analysis. *Can J Anaesth.* 2019;66(9):1095-1105.
27. Prestmo A, Hagen G, Sletvold O, et al. Comprehensive geriatric care for patients with hip fractures: a prospective, randomised, controlled trial. *Lancet.* 2015; 385(9978):1623-1633.
28. Swart E, Vasudeva BS, Makhni EC, et al. Dedicated perioperative hip fracture comanagement programs are cost-effective in high-volume centers: An economic analysis. *Clin Orthop Relat Res.* 2016 Jan;474(1):222-223.
29. American Academy of Orthopaedic Surgeons. Management of hip fractures in older adults: evidence-based clinical practice guideline. 2021. December 13, 2021. <https://www.aaos.org/hipfxcpq>.
30. van den Bekerom MPJ, Hilverdink EF, Sieverelt IN, et al. A comparison of hemiarthroplasty with total hip replacement for displaced intracapsular fracture of the femoral neck: a randomised controlled multicentre trial in patients aged 70 years and over. *Bone Joint Journal Brit.* 2010;92(10):1422-1428.
31. Bhandari M, Einhorn TA, Guyatt G, et al. Total Hip Arthroplasty or Hemiarthroplasty for Hip Fracture. *N Engl J Med.* 2019;381(23):2199-2208.
32. Baumgaertner M, Mai C, Oh J-K. Intertrochanteric Fractures, Nailing (short nail). *AO Surgery Reference*, Third edition (2022). AO Foundation. <https://surgeryreference.aofoundation.org/orthopedic-trauma/adult-trauma/proximal-femur/trochanteric-fracture-intertrochanteric/nailing-short-nail>. Accessed Aug 17, 2023
33. Pincus D, Ravi B, Wasserstein D, et al. Association between wait time and 30-day mortality in adults undergoing hip fracture surgery. *JAMA.* 2017 Nov 28;318(20):1994-2003.
34. Fu MC, Boddapati V, Gausden EB, et al. Surgery for a fracture of the hip within 24 hours of admission is independently associated with reduced short-term post-operative complications. *Bone Joint J.* 2017 Sep;99-B(9):1216-1222.
35. HIP ATTACK Investigators. Accelerated surgery versus standard care in hip fracture (HIP ATTACK): an international, randomised, controlled trial. *Lancet.* 2020;395(10225):698-708.
36. Neuman MD, Feng R, Carson JL, et al. Spinal Anesthesia or General Anesthesia for Hip Surgery in Older Adults. *N Engl J Med.* 2021;385(22):2025-2035.
37. Li T, Li J, Yuan L, et al. Effect of Regional vs General Anesthesia on Incidence of Postoperative Delirium in Older Patients Undergoing Hip Fracture Surgery: The RAGA Randomized Trial. *JAMA.* 2022;327(1):50-58.
38. Fleisher LA, Fleischmann KE, Auerbach AD, et al. 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery: a

report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2014;64:e77-137.

39. Grant PJ, Wesorick DH. Perioperative medicine for the hospitalized patient. *Med Clin N Am*. 2008;92:325-348.
40. Lee TH, Marcantonio ER, Mangione CM, et al. Derivation and prospective validation of a simple index for prediction of cardiac risk of major noncardiac surgery. *Circulation*. 1999;100:1043-1049.
41. Gupta PK, Gupta H, Sundaram A, et al. Development and validation of a risk calculator for prediction of cardiac risk after surgery. *Circulation*. 2011;124(4):381-387.
42. Glance LG, Faden E, Dutton RP, et al. Impact of the Choice of Risk Model for Identifying Low-risk Patients Using the 2014 American College of Cardiology/American Heart Association Perioperative Guidelines. *Anesthesiology*. 2018;129(5):889-900.
43. Cohn SL, Fernandez Ros N. Comparison of 4 Cardiac Risk Calculators in Predicting Postoperative Cardiac Complications After Noncardiac Operations. *Am J Cardiol*. 2018;121(1):125-130.
44. McFalls EO, Ward HB, Moritz TE, et al. Coronary artery revascularization before elective major vascular surgery. *N Engl J Med*. 2004;351:2795-804.
45. Stitgen A, Poludnianyk K, Dulaney-Crip E, et al. Adherence to preoperative cardiac clearance guidelines in hip fracture patients. *J Orthop Trauma*. 2015 Nov;29(11):500-3.
46. Canet J, Gallart L, Gomart C, et al. Prediction of postoperative pulmonary complications in a population-based surgical cohort. *Anesthesiology*. 2010;113(6):1338-50.
47. Smetana GW, Lawrence VA, Cornell JE. American College of Physicians. Preoperative pulmonary risk stratification for noncardiothoracic surgery: systematic review for the American College of Physicians. *Ann Intern Med*. 2006;144(8):581-95.
48. Lin H-S, Watts JN, Peel NM, et al. Frailty and post-operative outcomes in older surgical patients: a systematic review. *BMC Geriatr*. 2016 Aug 31;16(1):157.
49. Rockwood K, Song X, MacKnight C, et al. A global clinical measure of fitness and frailty in elderly people. *CMAJ*. 2005;173(5):489-95.
50. McIsaac DI, Taljaard M, Bryson GL, et al. Frailty as a predictor of death or new disability after surgery: A prospective cohort study. *Ann Surg*. 2018; Jul 24. doi: 10.1097/SLA.
51. Belga S, Majumdar SR, Kahlon S, et al. Comparing three different measures of frailty in medical inpatients: Multicenter prospective cohort study examining 30-day risk of readmission or death. *J Hosp Med*. 2016. August;11(8):556-562.

52. Berry SD, Rothbaum, RR, Kiel DP, et al. Association and clinical outcomes with surgical repair of hip fracture vs nonsurgical management in nursing home residents with advanced dementia. *JAMA Intern Med.* 2018;178(6):774-780.
53. Sullivan NM, Blake LE, George M, et al. Palliative care in the hip fracture patient. *Geriatr Orthop Surg Rehabil.* 2019;10:2151459319849801. doi:10.1177/2151459319849801.
54. Ko FC, Morrison RS. Hip fracture: a trigger for palliative care in vulnerable older adults. *JAMA Intern Med.* 2014;174(8):1281-1282
55. Geerts WH, Bergqvist D, Pineo GF, et al. Prevention of venous thromboembolism. American College of Chest Physicians Evidence-Based Clinical Practice Guidelines, 8th ed. *Chest.* 2008;133:381S-453S.
56. Pedersen AB, Ehrenstein V, Szépligeti SK, et al. Excess risk of venous thromboembolism in hip fracture patients and the prognostic impact of comorbidity. *Osteoporosis Int.* 2017;28(12):3421-3430.
57. Mula V, Parkh S, Sushesh S, et al. Venous thromboembolism rates after hip and knee arthroplasty and hip fracture. *BMC Musculoskel Disord.* 2020;21(1):95. doi:10.1186/s12891-020-3100-4.
58. Falck-Ytter Y, Francis CW, Johanson NA, et Al. Prevention of VTE in Orthopedic Surgery Patients. Antithrombotic Therapy and Prevention of Thrombosis, 9th ed. American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest.* 2012;141(2)(suppl):e278S-e325S.
59. Balk EM, Ellis AG, Di M, Adam GP, Trikalinos TA. Venous Thromboembolism Prophylaxis in Major Orthopedic Surgery: Systematic Review Update. Comparative Effectiveness Review No. 191. (Prepared by the Brown Evidence-based Practice Center under Contract No. 290-2015-00002-I.) AHRQ Publication No. 17-EHC021-EF. Rockville, MD: Agency for Healthcare Research and Quality; June 2017. www.effectivehealthcare.ahrq.gov/reports/final.cfm.
60. Muscatelli SR, Charter MA, Hallstrom BR. Time for an update? A look at current guidelines for venous thromboembolism prophylaxis after hip and knee arthroplasty and hip fracture. *Arthroplasty Today.* 2021;10:105-107.
61. Anderson DR, Morgano GP, Bennett C, et al. American Society of Hematology 2019 guidelines for management of venous thromboembolism: prevention of venous thromboembolism in surgical hospitalized patients. *Blood Adv.* 2019;3(23):3898-3944.
62. National Institute for Health Care and Clinical Excellence. Venous thromboembolism in over 16s: reducing the risk of hospital-acquired deep vein thrombosis or pulmonary embolism. NICE Guideline [NG89]. Aug 13, 2019. <https://www.nice.org.uk/guidance/ng89>

63. Goh, EL, Gurung PK, Ma S, et al. Direct oral anticoagulants in the prevention of venous thromboembolism following surgery for hip fracture in older adults: a population-based cohort study. *Geriatr Orthop Surg Rehab.* 2020;11:1-5.
64. Nederpelt CH, Bijman Q, Krijnen P, et al. Equivalence of DOACs and LMWH for thromboprophylaxis after hip fracture surgery: systemic review and meta-analysis. *Injury.* 2022;53:1169-1176.
65. Pulmonary Embolism Prevention (PEP) trial Collaborative Group. Prevention of pulmonary embolism and deep vein thrombosis with low dose aspirin: Pulmonary Embolism Prevention (PEP) trial. *Lancet.* 2000;355(9212):1295-1302.
66. Powers PJ, Gent M, Jay RM, et al. A randomized trial of less intense postoperative warfarin or aspirin therapy in the prevention of venous thromboembolism after surgery for fractured hip. *JAMA Intern Med.* 1989;149(3):771-774.
67. Drescher FS, Sirovich BE, Lee A, et al. Aspirin versus Anticoagulation for Prevention of Venous Thromboembolism Major Lower Extremity Orthopedic Surgery: A Systematic Review and Meta-Analysis. *J Hosp Med.* 2014;9:579-585.
68. Major Extremity Trauma Research Consortium. Aspirin or Low-Molecular-Weight heparin for thromboprophylaxis after a fracture. *N Engl J Med.* 2023;388(3):203-213.
69. Hung WW, Egol KA, Zuckerman JD, Siu AL. Hip fracture management tailoring Care for the older patient. *JAMA.* 2012;307(20):2185-2194.
70. Yang Y, Zhao X, Dong T, et al. Risk factors for postoperative delirium following hip fracture repair in elderly patients: a systematic review and meta-analysis. *Aging Clin Exp Res.* 2017 Apr;29(2):115-126.
71. Eamer G, Taheri A, Chen SS, et al. Comprehensive geriatric assessment for older people admitted to a surgical service. *Cochrane Database Syst Rev.* 2018 Jan 31;1:CD012485.
72. Shields L, Henderson V, Caslake R. Comprehensive geriatric assessment for prevention of delirium after hip fracture: a systematic review of randomized controlled trials. *J Am Geriatr Soc.* 2017 Jul;65(7):1559-1565.
73. The American Geriatrics Society Expert Panel on Postoperative Delirium in Older Adults. American Geriatrics Society Abstracted Clinical Practice Guideline for Postoperative Delirium in Older Adults. *J Am Geriatr Soc.* 2015 January;63(1):142-150.
74. Guay J, Kopp S. Peripheral nerve blocks for hip fractures in adults. *Cochrane Database Syst Rev.* 2020 Nov 25;11(11):CD001159.
75. Beaupre LA, Jones, CA, Saunders LD, et al. Best practices for elderly hip fracture patients. A systematic overview of the evidence. *J Gen Intern Med.* 2005;20:09-1025.

76. Baumgarten M, Margolis D, Berlin JA, et al. Risk factors for pressure ulcers among elderly hip fracture patients. *Wound Rep Reg*. 2003;11:96-103.
77. Chiari P, Forni C, Guberti M, et al. Predictive Factors for Pressure Ulcers in an Older Adult Population Hospitalized for Hip Fractures: A Prognostic Cohort Study. *PLoS One*. 2017;12(1): e0169909.
78. Hofman A, Geelkerken RH, Wille J, et al. Pressure sores and pressure-decreasing mattresses: controlled clinical trial. *Lancet*. 1994;343:568-571.
79. Siris ES, Adler R, Bilezikian J, et al. The clinical diagnosis of osteoporosis: a position statement from the National Bone Health Alliance Working Group. *Osteoporos Int*. 2014: 1439-1443.
80. Colon-Emeric C, Nordsletten L, Olson S, et al. Association between timing of zoledronic acid infusion and hip fracture healing. *Osteoporosis Int*. 2011;22:2329-2336.
81. LeBoff MS, Greenspan SL, Insogna KL, et al. The Clinician's Guide to Prevention and Treatment of Osteoporosis. *Osteoporosis Int*. 2022;33(10):2049-2102.
82. US Preventive Services Task Force. Vitamin D, calcium, or combined supplementation for the primary prevention of fractures in community-dwelling adults. US Preventive Services Task Force Recommendation Statement. *JAMA*. 2018;319(15):1592-1599.
83. Bolland MJ, Grey A, Avenell A. Effects of vitamin D supplementation on musculoskeletal health: a systematic review, meta-analysis, and trial sequential analysis. *Lancet Diabetes Endocrinol*. 2018 November;6(11):847-858.
84. Lyles KW, Colon-Emeric CS, Magaziner JS, et al. Zoledronic Acid in Reducing Clinical Fracture and Mortality after Hip Fracture. *N Engl J Med*. 2007;357:nihpa40967.
85. Li YT, Cai HF, Zhang ZL. Timing of the initiation of bisphosphonates after surgery for fracture healing: a systematic review and meta-analysis of randomized controlled trials. *Osteoporosis Int*. 2015;26:431-441.
86. Unnanuntana A, Laohaprasitiporn P, Jarusriwana A. Effect of bisphosphonate initiation at week 2 versus week 12 on short-term functional recovery after femoral neck fracture: a randomized controlled trial. *Arch Osteoporos*. 2017 Dec;12(1):27.
87. Desai RJ, Mahesri M, Abdia Y, et al. Association of Osteoporosis Medication Use after Hip Fracture with Prevention of Subsequent Nonvertebral Fractures An Instrumental Variable Analysis. *JAMA Network Open*. 2018;1(3):e180826.
88. Tan EK, Loh KP, Goff SL. Internal Medicine Hospitalists' Perceived Barriers and Recommendations for Optimizing Secondary Prevention of Osteoporotic Hip Fractures. *South Med J*. 2017 Dec;110(12):749-753.

89. Lv H, Yin P, Long A, et al. Clinical characteristics and risk factors of postoperative pneumonia after hip fracture surgery: a prospective cohort study. *Osteoporos Int.* 2016 Oct;27(10): 3001-9.